

Minimal Depth Vaginoplasty Post-Operative Instructions

General Instructions

- Someone must drive you home after your 3 day hospital stay and help you at home for 1-2 days.
- Get plenty of rest.
- Follow balanced diet with high protein and fluid intake.
- Decreased activity may promote constipation, so you may want to add more raw fruit to your diet and be sure to increase fluid intake.
- Take pain medication as prescribed. Do not take aspirin or any products containing aspirin unless approved by your surgeon.
- Do not drink alcohol when taking pain medications.
- Do not smoke, as smoking delays healing and increases the risk of complications.

Activities

- Start walking as soon as possible, this helps to reduce swelling and lowers the chance of blood clots
- Do not drive until you are no longer taking any pain medications (narcotics).
- Refrain from vigorous activity for 2-6 weeks. Increase activity gradually as tolerated.
- Avoid lifting anything over 5-10 pounds for 4 weeks.
- Resume social and employment activities in about 4 weeks (if not too strenuous).

Urination

- You may experience splashing and difficulty directing your urine stream for the first few months after surgery. This should resolve and become more streamline over time. If you continue to have urination stream difficulties, contact your surgeon.
- **If you experience the inability to or difficulty urinating after discharge from the hospital, contact your surgeon.**

Incision Care

- Oozing and bleeding from the incisions and vaginal pocket is expected. This can come and go for 3 to 6 months after surgery.
- Use feminine pads/gauze to soak up bleeding. Contact your doctor if you soak through 1 or more pad every hour.
- Wash surgical incisions and vaginal pocket gently with regular shower soap and water. No vaginal douching until cleared by your surgeon.
- No tub soaking/swimming for 1 month.

What to Expect

- Expect moderate drainage on your initial dressing, it is not uncommon for this to seep onto the surgical dressings.
- You may experience temporary soreness, bruising, swelling and tightness in the pubic area as well as discomfort in the incision area.
- Due to the location of your surgical site, slow wound healing and wound dehiscence that appears as 'raw flesh' is extremely common. This will heal with time, use bacitracin to the areas of slow healing 3 times daily.
- You may not have sensation in the pubic area, surrounding skin, or along the incisions. This usually returns with time but can take up to 12 to 18 months.
- You may have random, shooting pains and itching throughout your abdomen/back/flanks and along incisions for a few months. This is part of the normal healing process.

Appearance

- Most of the discoloration and swelling will subside in 4-6 weeks.
- Scars may be red and angry looking for 6 months. In time, these usually soften and fade.

Follow-Up Care

- Follow up within 1 week of surgery. Expect additional follow up appointments weekly, bi-weekly, and/or monthly post op as directed by your surgeon.
- All sutures will be dissolvable.

When to Call

- If you have severe or increased pain not relieved by medication.
- If you have any side effects to medications; such as, rash, nausea, headache, vomiting.
- If you have an oral temperature over 101F degrees.
- If you have sudden increased swelling or bruising.
- If you have increased redness and/or warmth along the incision.
- If you have bleeding from the incisions that is difficult to control with light pressure.

For Medical Questions, please contact: Katie Leemaster PA-C at katie@districtps.com or Dr. Praful Ramineni at doc@districtps.com

Your post op appointment is scheduled for _____